

Set Theory

集合论

杨睿之

yangruizhi@fudan.edu.cn

School of Philosophy, Fudan University

Fall 2015

Previously on Set Theory

Cofinality

\aleph_1

\aleph_1

- Set theoretical reals:

Baire Space ω^ω , subtree of $\omega^{<\omega}$

- Complexity of definable sets of reals:

- Borel sets

\aleph_1

- Projective sets

Analytic set Σ_1^1

Regularity Properties



Perfect set property

Definition

- A set $P \subset \omega^\omega$ is **perfect** if it is closed and has no isolated points, i.e. for each (basic) open set $U \subset \omega^\omega$ and $x \in P$, if $x \in U$, then $U \cap (P \setminus \{x\}) \neq \emptyset$
- A set $A \subset \omega^\omega$ has the perfect set property if it is either countable or contains a nonempty perfect subset

Perfect set property

Definition



- A set $P \subset \omega^\omega$ is **perfect** if it is closed and has no **isolated points**, i.e. for each (basic) open set $U \subset \omega^\omega$ and $x \in P$, if $x \in U$, then $U \cap (P \setminus \{x\}) \neq \emptyset$
- A set $A \subset \omega^\omega$ has the perfect set property if it is either countable or contains a nonempty perfect subset

Perfect set property

Definition

- A set $P \subset \omega^\omega$ is **perfect** if it is closed and has no **isolated points**, i.e. for each (basic) open set $U \subset \omega^\omega$ and $x \in P$, if $x \in U$, then $U \cap (P \setminus \{x\}) \neq \emptyset$
- A set $A \subset \omega^\omega$ has the **perfect set property** if it is either countable or contains a nonempty perfect subset

Perfect set property


Embed cantor space into a perfect set

$$2^\omega$$

Fact

$$\omega \vee 1$$

If P is perfect, then $\text{card } P = 2^{\aleph_0}$. Thus, if $A \subset \omega^\omega$ has the perfect set property, then A cannot be a witness of the failure of CH



$$f \langle 0 \rangle = s_0 \hat{\ } n_0$$

$$f \langle 1 \rangle = s_0 \hat{\ } m_0$$

$$f \langle 0, 0 \rangle = s_{00} \hat{\ } n_{00} \quad f \langle 1, 0 \rangle =$$

$$f \langle 0, 1 \rangle = s_{00} \hat{\ } m_{00}$$

Perfect set property

Theorem (Cantor-Bendixson)

Every closed set has the perfect set property

Let $C \subset \omega^\omega$ be closed.

Show that C is either countable or contains a nonempty perfect set

Perfect set property

Fact

Every analytic set has perfect set property

Perfect set property

Definition

A set $B \subset \omega^\omega$ is **Bernstein** if neither B nor $\omega^\omega \setminus B$ contains a perfect subset

Fact (AC)

There is a Bernstein set, which violates the perfect set property

Construct a Bernstein set

Perfect set property

Fact

- If $V = L$, then there is a \aleph_1^1 set of reals which does not have the perfect set property
- It follows from a large cardinal axiom that every projective set of reals has the perfect set property

Lesbesgue Measurable

For now, We consider the real line \mathbb{R}

Definition

Outer Measure

- For interval $I = (a, b)$, define the length of the interval $l(I) = b - a$
- For $A \subset \mathbb{R}$, the outer measure of A is

$$\mu^*(A) = \inf \left\{ \sum_{k \in \omega} l(I_k) \mid \langle I_k \rangle_{k \in \omega} \text{ is a open covering of } A \right\}$$

Lesbesgue Measurable

For now, We consider the real line \mathbb{R}

Definition

Outer Measure

- For interval $I = (a, b)$, define the length of the interval $l(I) = b - a$
- For $A \subset \mathbb{R}$, the **outer measure** of A is

$$\mu^*(A) = \inf \left\{ \sum_{k \in \omega} l(I_k) \mid \langle I_k \rangle_{k \in \omega} \text{ is a open covering of } A \right\}$$

Lesbesgue Measurable

Definition

Lesbesgue Measure

- A set $A \subset \mathbb{R}$ is (Lesbesgue) measurable if for every $X \subset \mathbb{R}$

$$\mu^*(A) = \mu^*(A \cap X) + \mu^*(A \setminus X)$$

- For Lesbesgue measurable set A , its Lesbesgue measure $\mu(A) = \mu^*(A)$

Lesbesgue Measurable

Fact

The collection of Lesbesgue measurable sets forms a σ -algebra. Thus every Borel set is Lesbesgue measurable

Theorem (Vitali)

There is a nonmeasurable set of reals

We find a Vitali set which is not measurable

Lesbesgue Measurable

Fact

Assuming $V = L$, there is a Σ_2^1 set of reals which is not measurable

Fact

It is provable from a large cardinal axiom that every projective set is measurable

Property of Baire

Definition

- A set $X \subset \omega^\omega$ is **nowhere dense** if its closure has empty interior, i.e. no nonempty open set is contained in its closure
- A set $M \subset \omega^\omega$ is **meager** if it is the union of countably many nowhere dense sets
- A set $A \subset \omega^\omega$ is said to have the **property of Baire** if it differs from an open set by a meager set

Property of Baire

Fact

The family of sets with the property of Baire forms a σ -algebra, which follows that every Borel set has the property of Baire

Fact

Neither Bernstein set nor Vitali set has the property of Baire

Property of Baire

Fact

The family of sets with the property of Baire forms a σ -algebra, which follows that every Borel set has the property of Baire

Fact

Neither Bernstein set nor Vitali set has the property of Baire

Determinacy

Definition (The game on ω)

Let $A \subset \omega^\omega$. We define the game G_A as follow

- A **play** is a sequence $\langle a_0, b_0, a_1, b_1, \dots \rangle$ of natural numbers, where $\langle a_0, a_1, \dots \rangle$ is the **play of player I** and $\langle b_0, b_1, \dots \rangle$ is the **play of player II**
- **Player I wins** the play if $\langle a_0, b_0, a_1, b_1, \dots \rangle \in A$

Determinacy

Definition (The game on ω)

- A **strategy** is a function $\sigma : \omega^{<\omega} \rightarrow \omega$. Given $b = \langle b_0, b_1, \dots \rangle$, $\sigma * b = \langle \sigma(\emptyset), b_0, \sigma(\langle \sigma(\emptyset), b_0 \rangle), b_1, \dots \rangle$ is the play of the player I's strategy σ against b . Similar is defined for player II's $a * \sigma$
- A **winning strategy for player I** is a strategy σ such that $\sigma * b \in A$ for all $b \in \omega^\omega$
- A **winning strategy for player II** is a strategy σ such that $a * \sigma \notin A$ for all $a \in \omega^\omega$

Determinacy

Definition

Axiom of determinacy

- A set $A \subset \omega^\omega$ is **determined** if either player I or player II has a winning strategy in G_A
- Axiom of determinacy (AD): Every subset $A \subset \omega^\omega$ is determined

Determinacy

Fact

Π_n^1 -determinacy implies that every Σ_{n+1}^1 set is Lebesgue measurable, has the property of Baire and the perfect set property

Corollary (AC)

\neg AD

Determinacy

Fact

Π_n^1 -determinacy implies that every Σ_{n+1}^1 set is Lebesgue measurable, has the property of Baire and the perfect set property

Corollary (AC)

\neg AD

Determinacy

Fact

Π_n^1 -determinacy implies that every Σ_{n+1}^1 set is Lebesgue measurable, has the property of Baire and the perfect set property

Corollary

If every projective set is determined (**PD**), then every projective set has the regularity properties

Determinacy

Theorem

Martin-Steel The existence of n Woodin cardinal with a measurable cardinal above them all imply Π_{n+1}^1 -determinacy. Therefore the existence of infinitely many Woodin cardinals implies PD

Next on Set Theory

- Constructible sets (Gödel's L)
- Forcing
- More on large cardinals and descriptive set theory