

# Equivalence relations and Borel reduction

2016 Chinese Mathematical Logic Conference  
Fudan University

Longyun Ding

School of Mathematical Sciences  
Nankai University

22 May 2016

# Outline

- 1 Equivalence relations and invariants
- 2 Dichotomy theorems
- 3 Orbit equivalence relations
- 4  $\Sigma_1^1$  equivalence relations

# Classical examples (1)

## Example

For  $A, B \in \mathbb{C}^{m \times n}$ , define  $A \sim B \iff A = TBS$ , where  $T, S$  are invertible matrices.

Let  $r(A)$  be rank of  $A$ . Then

$$A \sim B \iff r(A) = r(B).$$

## Example

For  $A, B \in \mathbb{C}^{n \times n}$ , define  $A \approx B \iff A = TBT^{-1}$ , where  $T$  is an invertible matrix.

Let  $J(A)$  be the Jordan normal form of  $A$ . Then

$$A \approx B \iff J(A) = J(B).$$

## Classical examples (1)

### Example

For  $A, B \in \mathbb{C}^{m \times n}$ , define  $A \sim B \iff A = TBS$ , where  $T, S$  are invertible matrices.

Let  $r(A)$  be rank of  $A$ . Then

$$A \sim B \iff r(A) = r(B).$$

### Example

For  $A, B \in \mathbb{C}^{n \times n}$ , define  $A \approx B \iff A = TBT^{-1}$ , where  $T$  is an invertible matrix.

Let  $J(A)$  be the Jordan normal form of  $A$ . Then

$$A \approx B \iff J(A) = J(B).$$

## Classical examples (2)

### Example

Every finitely generated abelian group  $G$  is isomorphic to a direct sum

$$\mathbb{Z}^m \oplus \bigoplus_{i=0}^n \bigoplus_{j=0}^{e_i} \mathbb{Z}(p_i^j)^{t_{ij}}.$$

Let  $M(G) = (m, t_{ij})_{i \leq n, j \leq e_i}$ . Then

$$G \cong H \iff M(G) = M(H).$$

**Note:**  $r(A)$ ,  $J(A)$ ,  $M(G)$  are not continuous mappings.

## Classical examples (3)

### Example

For compact topological spaces  $X$ , denote  $C(X)$  the spaces of all continuous function  $X \rightarrow \mathbb{C}$  equipped with the sup norm.

### Theorem (Gelfand-Naimark)

*Let  $X, Y$  be compact Hausdorff spaces. Then  $X$  is homeomorphic to  $Y$  iff  $C(X)$  is isomorphic to  $C(Y)$  (as a  $C^*$ -algebra).*

## Classical examples (3)

### Example

For compact topological spaces  $X$ , denote  $C(X)$  the spaces of all continuous function  $X \rightarrow \mathbb{C}$  equipped with the sup norm.

### Theorem (Gelfand-Naimark)

*Let  $X, Y$  be compact Hausdorff spaces. Then  $X$  is homeomorphic to  $Y$  iff  $C(X)$  is isomorphic to  $C(Y)$  (as a  $C^*$ -algebra).*

# Reduction

## Definition

Let  $E, F$  be two equivalence relations on  $X, Y$  respectively,  $\theta : X \rightarrow Y$  is a **reduction** of  $E$  to  $F$  if

$$aEb \iff \theta(a)F\theta(b)$$

for  $a, b \in X$ .

## Fact

Let  $f : X/E \rightarrow X$  be a choice function, and let  $\theta(a) = f([a]_E)$ . Then  $\theta$  is a reduction of  $E$  to  $\text{id}(X)$ .

**Note:** We need some restrictions on reduction mapping!



# Reduction

## Definition

Let  $E, F$  be two equivalence relations on  $X, Y$  respectively,  $\theta : X \rightarrow Y$  is a **reduction** of  $E$  to  $F$  if

$$aEb \iff \theta(a)F\theta(b)$$

for  $a, b \in X$ .

## Fact

Let  $f : X/E \rightarrow X$  be a choice function, and let  $\theta(a) = f([a]_E)$ . Then  $\theta$  is a reduction of  $E$  to  $\text{id}(X)$ .

**Note:** We need some restrictions on reduction mapping!

# Polish spaces

## Definition

**Polish space:** a separable, completely metrizable topological space.

## Example

- 1  $\mathbb{C}^{m \times n}; \mathbb{R}^n$ ; separable Banach spaces;
- 2  $\mathcal{C} = \{0, 1\}^{\mathbb{N}}; \mathcal{N} = \mathbb{N}^{\mathbb{N}}$ ;
- 3 For a countable abelian group  $G$ , note that

$$R_G = \{(a, b, c) : a = b + c\} \subseteq \mathbb{N}^3.$$

“Countable abelian groups” is a closed subset of  $\{0, 1\}^{\mathbb{N}^3} \cong \mathcal{C}$ .

# Polish spaces

## Definition

**Polish space:** a separable, completely metrizable topological space.

## Example

- 1  $\mathbb{C}^{m \times n}$ ;  $\mathbb{R}^n$ ; separable Banach spaces;
- 2  $\mathcal{C} = \{0, 1\}^{\mathbb{N}}$ ;  $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$ ;
- 3 For a countable abelian group  $G$ , note that

$$R_G = \{(a, b, c) : a = b + c\} \subseteq \mathbb{N}^3.$$

“Countable abelian groups” is a closed subset of  $\{0, 1\}^{\mathbb{N}^3} \cong \mathcal{C}$ .

# Polish spaces

## Definition

**Polish space:** a separable, completely metrizable topological space.

## Example

- 1  $\mathbb{C}^{m \times n}$ ;  $\mathbb{R}^n$ ; separable Banach spaces;
- 2  $\mathcal{C} = \{0, 1\}^{\mathbb{N}}$ ;  $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$ ;
- 3 For a countable abelian group  $G$ , note that

$$R_G = \{(a, b, c) : a = b + c\} \subseteq \mathbb{N}^3.$$

“Countable abelian groups” is a closed subset of  $\{0, 1\}^{\mathbb{N}^3} \cong \mathcal{C}$ .

# Polish spaces

## Definition

**Polish space:** a separable, completely metrizable topological space.

## Example

- 1  $\mathbb{C}^{m \times n}$ ;  $\mathbb{R}^n$ ; separable Banach spaces;
- 2  $\mathcal{C} = \{0, 1\}^{\mathbb{N}}$ ;  $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$ ;
- 3 For a countable abelian group  $G$ , note that

$$R_G = \{(a, b, c) : a = b + c\} \subseteq \mathbb{N}^3.$$

“Countable abelian groups” is a closed subset of  $\{0, 1\}^{\mathbb{N}^3} \cong \mathcal{C}$ .

## Borel sets, Borel functions and Borel reductions

### Definition

$\mathbf{B}(X)$ : **Borel sets** of  $X$  is the  $\sigma$ -algebra generated by open sets.

$\mathbf{B}(X)$  contains all open, closed,  $F_\sigma, G_\delta, G_{\delta\sigma}, F_{\sigma\delta}$ , etc., sets.

Let  $X, Y$  be two Polish spaces.

### Definition

A function  $f : X \rightarrow Y$  is **Borel function** if  $f^{-1}(U)$  is Borel for  $U$  open in  $Y$ .

Let  $E, F$  be equivalence relations on  $X, Y$  respectively.

$E \leq_B F$ : There is a Borel reduction of  $E$  to  $F$ ;

$E \sim_B F$ :  $E \leq_B F$  and  $F \leq_B E$ ;

$E <_B F$ :  $E \leq_B F$  but not  $F \leq_B E$ .

## Borel sets, Borel functions and Borel reductions

### Definition

$\mathbf{B}(X)$ : **Borel sets** of  $X$  is the  $\sigma$ -algebra generated by open sets.

$\mathbf{B}(X)$  contains all open, closed,  $F_\sigma, G_\delta, G_{\delta\sigma}, F_{\sigma\delta}$ , etc., sets.

Let  $X, Y$  be two Polish spaces.

### Definition

A function  $f : X \rightarrow Y$  is **Borel function** if  $f^{-1}(U)$  is Borel for  $U$  open in  $Y$ .

Let  $E, F$  be equivalence relations on  $X, Y$  respectively.

$E \leq_B F$ : There is a Borel reduction of  $E$  to  $F$ ;

$E \sim_B F$ :  $E \leq_B F$  and  $F \leq_B E$ ;

$E <_B F$ :  $E \leq_B F$  but not  $F \leq_B E$ .

# Borel sets, Borel functions and Borel reductions

## Definition

$\mathbf{B}(X)$ : **Borel sets** of  $X$  is the  $\sigma$ -algebra generated by open sets.

$\mathbf{B}(X)$  contains all open, closed,  $F_\sigma, G_\delta, G_{\delta\sigma}, F_{\sigma\delta}$ , etc., sets.

Let  $X, Y$  be two Polish spaces.

## Definition

A function  $f : X \rightarrow Y$  is **Borel function** if  $f^{-1}(U)$  is Borel for  $U$  open in  $Y$ .

Let  $E, F$  be equivalence relations on  $X, Y$  respectively.

$E \leq_B F$ : There is a Borel reduction of  $E$  to  $F$ ;

$E \sim_B F$ :  $E \leq_B F$  and  $F \leq_B E$ ;

$E <_B F$ :  $E \leq_B F$  but not  $F \leq_B E$ .



# Standard Borel spaces

## Definition

A measurable space  $(X, \mathcal{S})$  is a *standard Borel space* if there is a Polish topology  $\tau$  on  $X$  with  $\mathcal{S} = \mathbf{B}(X, \tau)$ .

## Theorem

*Let  $X$  be a Polish space,  $Y \subseteq X$ . Then  $(Y, \mathbf{B}(Y))$  is a standard Borel space iff  $Y$  is a Borel subset of  $X$ .*

## Example

“finitely generated abelian groups” is a Borel subset of “countable abelian groups”, so it is a standard Borel space.

# Standard Borel spaces

## Definition

A measurable space  $(X, \mathcal{S})$  is a *standard Borel space* if there is a Polish topology  $\tau$  on  $X$  with  $\mathcal{S} = \mathbf{B}(X, \tau)$ .

## Theorem

*Let  $X$  be a Polish space,  $Y \subseteq X$ . Then  $(Y, \mathbf{B}(Y))$  is a standard Borel space iff  $Y$  is a Borel subset of  $X$ .*

## Example

“finitely generated abelian groups” is a Borel subset of “countable abelian groups”, so it is a standard Borel space.

# Standard Borel spaces

## Definition

A measurable space  $(X, \mathcal{S})$  is a *standard Borel space* if there is a Polish topology  $\tau$  on  $X$  with  $\mathcal{S} = \mathbf{B}(X, \tau)$ .

## Theorem

*Let  $X$  be a Polish space,  $Y \subseteq X$ . Then  $(Y, \mathbf{B}(Y))$  is a standard Borel space iff  $Y$  is a Borel subset of  $X$ .*

## Example

“finitely generated abelian groups” is a Borel subset of “countable abelian groups”, so it is a standard Borel space.

## Effros Borel spaces

### Example (Effros Borel spaces)

Given a Polish space  $X$ , we denote by  $F(X)$  the set of closed subsets of  $X$ . We endow  $F(X)$  with the  $\sigma$ -algebra generated by the sets

$$\{F \in F(X) : F \cap U \neq \emptyset\},$$

where  $U$  varies over open subsets of  $X$ .

### Fact

*If  $X$  is Polish, the Effros Borel space of  $F(X)$  is a standard Borel space.*

# About Gelfand-Naimark's theorem

## Fact

- ① *Every compact metric space homeomorphic to a closed subset of Hilbert Cub  $[0, 1]^{\mathbb{N}}$ .*
- ② *Every separable Banach space isometrically isomorphic to a closed linear subspace of  $C[0, 1]$ .*

Let  $\text{Hom}_{\text{cpt}}$  be the homeomorphism relation on  $F([0, 1]^{\mathbb{N}})$ , and let  $\cong_{\text{SB}}$  be the isometrically isomorphism on  $\text{Subs}(C[0, 1]) \subseteq F(C[0, 1])$  of all closed linear subspaces of  $C[0, 1]$ . Then

$$\text{Hom}_{\text{cpt}} \leq_B \cong_{\text{SB}} .$$

# About Gelfand-Naimark's theorem

## Fact

- ① *Every compact metric space homeomorphic to a closed subset of Hilbert Cub  $[0, 1]^{\mathbb{N}}$ .*
- ② *Every separable Banach space isometrically isomorphic to a closed linear subspace of  $C[0, 1]$ .*

Let  $\text{Hom}_{\text{cpt}}$  be the homeomorphism relation on  $F([0, 1]^{\mathbb{N}})$ , and let  $\cong_{\text{SB}}$  be the isometrically isomorphism on  $\text{Subs}(C[0, 1]) \subseteq F(C[0, 1])$  of all closed linear subspaces of  $C[0, 1]$ . Then

$$\text{Hom}_{\text{cpt}} \leq_B \cong_{\text{SB}} .$$

# About Gelfand-Naimark's theorem

## Fact

- ① *Every compact metric space homeomorphic to a closed subset of Hilbert Cub  $[0, 1]^{\mathbb{N}}$ .*
- ② *Every separable Banach space isometrically isomorphic to a closed linear subspace of  $C[0, 1]$ .*

Let  $\text{Hom}_{\text{cpt}}$  be the homeomorphism relation on  $F([0, 1]^{\mathbb{N}})$ , and let  $\cong_{\text{SB}}$  be the isometrically isomorphism on  $\text{Subs}(C[0, 1]) \subseteq F(C[0, 1])$  of all closed linear subspaces of  $C[0, 1]$ . Then

$$\text{Hom}_{\text{cpt}} \leq_B \cong_{\text{SB}} .$$

# About Gelfand-Naimark's theorem

## Fact

- ① *Every compact metric space homeomorphic to a closed subset of Hilbert Cub  $[0, 1]^{\mathbb{N}}$ .*
- ② *Every separable Banach space isometrically isomorphic to a closed linear subspace of  $C[0, 1]$ .*

Let  $\text{Hom}_{\text{cpt}}$  be the homeomorphism relation on  $F([0, 1]^{\mathbb{N}})$ , and let  $\cong_{\text{SB}}$  be the isometrically isomorphism on  $\text{Subs}(C[0, 1]) \subseteq F(C[0, 1])$  of all closed linear subspaces of  $C[0, 1]$ .

Then

$$\text{Hom}_{\text{cpt}} \leq_B \cong_{\text{SB}} .$$



# About Gelfand-Naimark's theorem

## Fact

- ① *Every compact metric space homeomorphic to a closed subset of Hilbert Cub  $[0, 1]^{\mathbb{N}}$ .*
- ② *Every separable Banach space isometrically isomorphic to a closed linear subspace of  $C[0, 1]$ .*

Let  $\text{Hom}_{\text{cpt}}$  be the homeomorphism relation on  $F([0, 1]^{\mathbb{N}})$ , and let  $\cong_{\text{SB}}$  be the isometrically isomorphism on  $\text{Subs}(C[0, 1]) \subseteq F(C[0, 1])$  of all closed linear subspaces of  $C[0, 1]$ . Then

$$\text{Hom}_{\text{cpt}} \leq_B \cong_{\text{SB}} .$$

## Smooth equivalence relations

We denote by  $\text{id}(X)$  the *identity relation* on  $X$ .

$$\text{id}(\mathbb{N}) <_B \text{id}(\mathbb{N}) <_B \text{id}(\mathbb{R}).$$

### Definition

We say  $E$  is **smooth** if  $E \leq_B \text{id}(\mathbb{R})$ .

### Fact

Let  $X, Y$  be Polish spaces, then  $X$  is Borel isomorphic to  $Y$  (i.e., there is a Borel bijection from  $X$  to  $Y$ ) iff  $|X| = |Y|$ .

## Smooth equivalence relations

We denote by  $\text{id}(X)$  the *identity relation* on  $X$ .

$$\text{id}(n) <_B \text{id}(\mathbb{N}) <_B \text{id}(\mathbb{R}).$$

### Definition

We say  $E$  is **smooth** if  $E \leq_B \text{id}(\mathbb{R})$ .

### Fact

*Let  $X, Y$  be Polish spaces, then  $X$  is Borel isomorphic to  $Y$  (i.e., there is a Borel bijection from  $X$  to  $Y$ ) iff  $|X| = |Y|$ .*

## Smooth equivalence relations

We denote by  $\text{id}(X)$  the *identity relation* on  $X$ .

$$\text{id}(n) <_B \text{id}(\mathbb{N}) <_B \text{id}(\mathbb{R}).$$

### Definition

We say  $E$  is **smooth** if  $E \leq_B \text{id}(\mathbb{R})$ .

### Fact

Let  $X, Y$  be Polish spaces, then  $X$  is Borel isomorphic to  $Y$  (i.e., there is a Borel bijection from  $X$  to  $Y$ ) iff  $|X| = |Y|$ .

## $\Sigma_1^1$ sets and $\Pi_1^1$ sets

### Definition

Let  $X$  be a Polish space. A subset  $A \subseteq X$  is **analytic** (or  $\Sigma_1^1$ ) if there is a Polish space  $Y$  and a closed subset  $C \subseteq X \times Y$  such that

$$x \in A \iff \exists y \in Y ((x, y) \in C).$$

A subset  $A \subseteq X$  is **co-analytic** (or  $\Pi_1^1$ ) if  $X \setminus A$  is  $\Sigma_1^1$ .

### Theorem (Suslin)

*Let  $A \subseteq X$ . Then  $A$  is Borel iff it is both  $\Sigma_1^1$  and  $\Pi_1^1$ .*

**fact:** All  $\sigma(\Sigma_1^1)$  sets in  $\mathbb{R}$  are Lebesgue measurable.

## $\Sigma_1^1$ sets and $\Pi_1^1$ sets

### Definition

Let  $X$  be a Polish space. A subset  $A \subseteq X$  is **analytic** (or  $\Sigma_1^1$ ) if there is a Polish space  $Y$  and a closed subset  $C \subseteq X \times Y$  such that

$$x \in A \iff \exists y \in Y ((x, y) \in C).$$

A subset  $A \subseteq X$  is **co-analytic** (or  $\Pi_1^1$ ) if  $X \setminus A$  is  $\Sigma_1^1$ .

### Theorem (Suslin)

*Let  $A \subseteq X$ . Then  $A$  is Borel iff it is both  $\Sigma_1^1$  and  $\Pi_1^1$ .*

**fact:** All  $\sigma(\Sigma_1^1)$  sets in  $\mathbb{R}$  are Lebesgue measurable.

$\Sigma_1^1$  sets and  $\Pi_1^1$  sets

## Definition

Let  $X$  be a Polish space. A subset  $A \subseteq X$  is **analytic** (or  $\Sigma_1^1$ ) if there is a Polish space  $Y$  and a closed subset  $C \subseteq X \times Y$  such that

$$x \in A \iff \exists y \in Y ((x, y) \in C).$$

A subset  $A \subseteq X$  is **co-analytic** (or  $\Pi_1^1$ ) if  $X \setminus A$  is  $\Sigma_1^1$ .

## Theorem (Suslin)

*Let  $A \subseteq X$ . Then  $A$  is Borel iff it is both  $\Sigma_1^1$  and  $\Pi_1^1$ .*

**fact:** All  $\sigma(\Sigma_1^1)$  sets in  $\mathbb{R}$  are Lebesgue measurable.

# Non-smooth equivalence relations

$T$  is a **transversal** for  $E$  if  $T$  meets each  $E$ -class at exactly one point.

## Theorem

let  $E$  be a Borel equivalence relation on a Polish space  $X$ . If  $E$  is smooth, then  $E$  has a  $\sigma(\Sigma_1^1)$  transversal.

## Definition

**Vitali equivalence relation:** For  $x, y \in \mathbb{R}$  we define

$$xE_vy \iff x - y \in \mathbb{Q}.$$

**Note:** Any transversal of  $E_v$  is not Lebesgue measurable, so  $E_v$  is not smooth.



## Non-smooth equivalence relations

$T$  is a **transversal** for  $E$  if  $T$  meets each  $E$ -class at exactly one point.

### Theorem

let  $E$  be a Borel equivalence relation on a Polish space  $X$ . If  $E$  is smooth, then  $E$  has a  $\sigma(\Sigma_1^1)$  transversal.

### Definition

**Vitali equivalence relation:** For  $x, y \in \mathbb{R}$  we define

$$xE_vy \iff x - y \in \mathbb{Q}.$$

**Note:** Any transversal of  $E_v$  is not Lebesgue measurable, so  $E_v$  is not smooth.

# Outline

- 1 Equivalence relations and invariants
- 2 Dichotomy theorems**
- 3 Orbit equivalence relations
- 4  $\Sigma_1^1$  equivalence relations


# 1st dichotomy theorem

We say an equivalence relation  $E$  on  $X$  is Borel,  $\Sigma_1^1$ , or  $\Pi_1^1$  if it is so in  $X^2$ .

## Theorem (Silver, 1980)

*Let  $E$  be a  $\Pi_1^1$  equivalence relation. Then  $E$  has either at most countably many or perfectly many equivalence classes, i.e.,*

$$E \leq_B \text{id}(\mathbb{N}) \text{ or } \text{id}(\mathbb{R}) \leq_B E.$$



## 2nd dichotomy theorem

### Definition

$E_0$  is the equivalence relation on  $2^{\mathbb{N}}$  defined by

$$xE_0y \iff \exists m \forall n \geq m (x(n) = y(n)).$$

**Fact:**  $E_0 \sim_B E_v = \mathbb{R}/\mathbb{Q}$ .

Theorem (Harrington-Kechris-Louveau, 1990)

*Let  $E$  be a Borel equivalence relation. Then either  $E \leq_B \text{id}(\mathbb{R})$  or  $E_0 \leq_B E$ .*

## 2nd dichotomy theorem

### Definition


$E_0$  is the equivalence relation on  $2^{\mathbb{N}}$  defined by

$$xE_0y \iff \exists m \forall n \geq m (x(n) = y(n)).$$

**Fact:**  $E_0 \sim_B E_v = \mathbb{R}/\mathbb{Q}$ .

### Theorem (Harrington-Kechris-Louveau, 1990)

Let  $E$  be a Borel equivalence relation. Then either  $E \leq_B \text{id}(\mathbb{R})$  or  $E_0 \leq_B E$ .



## 3rd dichotomy theorem

### Definition

$E_1$  is the equivalence relation on  $\mathbb{R}^{\mathbb{N}}$  defined by

$$xE_1y \iff \exists m \forall n \geq m (x(n) = y(n)).$$

**Fact:**  $E_1 = \mathbb{R}^{\mathbb{N}}/c_{00}$ , where  $c_{00} = \bigcup_n \mathbb{R}^n$ .

Theorem (Kechris-Louveau, 1997)

*If  $E \leq_B E_1$ , then  $E \leq_B E_0$  or  $E \sim_B E_1$ .*



## 3rd dichotomy theorem

### Definition


$E_1$  is the equivalence relation on  $\mathbb{R}^{\mathbb{N}}$  defined by

$$xE_1y \iff \exists m \forall n \geq m (x(n) = y(n)).$$

**Fact:**  $E_1 = \mathbb{R}^{\mathbb{N}}/c_{00}$ , where  $c_{00} = \bigcup_n \mathbb{R}^n$ .

### Theorem (Kechris-Louveau, 1997)

If  $E \leq_B E_1$ , then  $E \leq_B E_0$  or  $E \sim_B E_1$ .



# 4th dichotomy theorem

## Definition

Let  $E$  be an equivalence relation on  $X$ . The equivalence relation  $E^\omega$  on  $X^\mathbb{N}$  defined by

$$xE^\omega y \iff \forall n(x(n)Ey(n)).$$

**Fact:**  $E_0^\omega \sim_B \mathbb{R}^\mathbb{N}/\mathbb{Q}^\mathbb{N}$ .

Theorem (Hjorth-Kechris, 1997)

*If  $E \leq_B E_0^\omega$ , then  $E \leq_B E_0$  or  $E \sim_B E_0^\omega$ .*

## 4th dichotomy theorem

### Definition


Let  $E$  be an equivalence relation on  $X$ . The equivalence relation  $E^\omega$  on  $X^\mathbb{N}$  defined by

$$xE^\omega y \iff \forall n(x(n)Ey(n)).$$

**Fact:**  $E_0^\omega \sim_B \mathbb{R}^\mathbb{N}/\mathbb{Q}^\mathbb{N}$ .

### Theorem (Hjorth-Kechris, 1997)

If  $E \leq_B E_0^\omega$ , then  $E \leq_B E_0$  or  $E \sim_B E_0^\omega$ .



# Outline

- 1 Equivalence relations and invariants
- 2 Dichotomy theorems
- 3 Orbit equivalence relations**
- 4  $\Sigma_1^1$  equivalence relations

# Polish $G$ -spaces and orbit equivalence relations

## Definition

**Polish group:** A topological group whose underlying space is Polish.

$G$ : Polish group,

$X$ : Polish space,

$a : G \times X \rightarrow X$ : continuous  $G$ -action on  $X$ .

## Definition

**Orbit equivalence relation:**

$$xE_G^X y \iff \exists g \in G (g \cdot x = y).$$

Any  $E_G^X$  is  $\Sigma_1^1$  equivalence relation.

## Polish $G$ -spaces and orbit equivalence relations

### Definition

**Polish group:** A topological group whose underlying space is Polish.

$G$ : Polish group,

$X$ : Polish space,

$a : G \times X \rightarrow X$ : continuous  $G$ -action on  $X$ .

### Definition

**Orbit equivalence relation:**

$$xE_G^X y \iff \exists g \in G (g \cdot x = y).$$

Any  $E_G^X$  is  $\Sigma_1^1$  equivalence relation.



# Polish $G$ -spaces and orbit equivalence relations

## Definition

**Polish group:** A topological group whose underlying space is Polish.

$G$ : Polish group,

$X$ : Polish space,

$a : G \times X \rightarrow X$ : continuous  $G$ -action on  $X$ .

## Definition

**Orbit equivalence relation:**

$$xE_G^X y \iff \exists g \in G (g \cdot x = y).$$

Any  $E_G^X$  is  $\Sigma_1^1$  equivalence relation.

# Polish groups

- 1 Countable groups with discrete topology.
- 2  $S_\infty$ : all bijections of  $\mathbb{N} \rightarrow \mathbb{N}$  with pointwise convergence topology.
- 3 **Universal Polish group**  $H([0, 1]^\mathbb{N})$ : every Polish group is isomorphic to one of its closed subgroups. (Uspenskii, 1986)
- 4 **Surjectively universal Polish group**  $\overline{F}_\Gamma(\mathcal{N}_\omega)$ : every Polish group is isomorphic to one of its topological quotient group. (D. 2012)

# Polish groups

- 1 Countable groups with discrete topology.
- 2  $S_\infty$ : all bijections of  $\mathbb{N} \rightarrow \mathbb{N}$  with pointwise convergence topology.
- 3 **Universal Polish group**  $H([0, 1]^\mathbb{N})$ : every Polish group is isomorphic to one of its closed subgroups. (Uspenskii, 1986)
- 4 **Surjectively universal Polish group**  $\overline{F}_\Gamma(\mathcal{N}_\omega)$ : every Polish group is isomorphic to one of its topological quotient group. (D. 2012)

# Polish groups

- 1 Countable groups with discrete topology.
- 2  $S_\infty$ : all bijections of  $\mathbb{N} \rightarrow \mathbb{N}$  with pointwise convergence topology.
- 3 **Universal Polish group**  $H([0, 1]^{\mathbb{N}})$ : every Polish group is isomorphic to one of its closed subgroups. (Uspenskii, 1986)
- 4 **Surjectively universal Polish group**  $\overline{F}_\Gamma(\mathcal{N}_\omega)$ : every Polish group is isomorphic to one of its topological quotient group. (D. 2012)

# Polish groups

- 1 Countable groups with discrete topology.
- 2  $S_\infty$ : all bijections of  $\mathbb{N} \rightarrow \mathbb{N}$  with pointwise convergence topology.
- 3 **Universal Polish group**  $H([0, 1]^{\mathbb{N}})$ : every Polish group is isomorphic to one of its closed subgroups. (Uspenskii, 1986)
- 4 **Surjectively universal Polish group**  $\overline{F}_\Gamma(\mathcal{N}_\omega)$ : every Polish group is isomorphic to one of its topological quotient group. (D. 2012)


# Countable Borel equivalence relations

An equivalence relation  $E$  on  $X$  is countable if every equivalence class of  $E$  is countable.

## Theorem (Feldman - Moore)

*Let  $E$  be a countable Borel equivalence relation on a Polish space  $X$ . Then  $E = E_G^X$  for some countable discrete group  $G$ .*

$E_\infty$ : an universal countable Borel equivalence relation.



## Below $E_0$

Theorem (Jackson-Kechris-Louveau, 2002)

$$E_{\mathbb{R}^n}^X \leq_B E_0.$$

Theorem (Gao-Jackson, 2015)

*For any countable abelian discrete group  $G$ ,  $E_G^X \leq_B E_0$ .*

Theorem (D.-Gao, 2015)

*Let  $G$  be a abelian closed subgroup of  $S_\infty$  and  $E_G^X \leq E_\infty$ , then  $E_G^X \leq_B E_0$ .*



## Below $E_0$

Theorem (Jackson-Kechris-Louveau, 2002)

$$E_{\mathbb{R}^n}^X \leq_B E_0.$$

Theorem (Gao-Jackson, 2015)

*For any countable abelian discrete group  $G$ ,  $E_G^X \leq_B E_0$ .*

Theorem (D.-Gao, 2015)

*Let  $G$  be a abelian closed subgroup of  $S_\infty$  and  $E_G^X \leq E_\infty$ , then  $E_G^X \leq_B E_0$ .*

## Below $E_0$

Theorem (Jackson-Kechris-Louveau, 2002)

$$E_{\mathbb{R}^n}^X \leq_B E_0.$$

Theorem (Gao-Jackson, 2015)

*For any countable abelian discrete group  $G$ ,  $E_G^X \leq_B E_0$ .*

Theorem (D.-Gao, 2015)

*Let  $G$  be a abelian closed subgroup of  $S_\infty$  and  $E_G^X \leq E_\infty$ , then  $E_G^X \leq_B E_0$ .*


## $S_\infty$ -spaces

**Borel complete**  $E_{S_\infty}^\infty$ : every  $E_{S_\infty}^X$  is Borel reducible to it.

### Theorem

*The isomorphism relations of all countable graphs, countable trees, countable linear orderings and countable groups are Borel complete, i.e.,*

$$E_{S_\infty}^\infty \sim_B (\cong_{\text{Graph}}) \sim_B (\cong_{\text{Tr}}) \sim_B (\cong_{\text{LO}}) \sim_B (\cong_{\text{Group}}).$$



# Upper bound of $E_G^X$

## Theorem (Gao-Kechris, 2003)

Let  $\cong_{\text{PM}}$  be the isometric isomorphism relation between Polish metric spaces. Then  $E_G^X \leq_B \cong_{\text{PM}}$  for every Polish  $G$ -space  $X$ .

## Theorem (Melleray-Weaver, 2007)

$\cong_{\text{PM}} \sim \cong_{\text{SB}}$ .

## Theorem (Zielinski, 2015)

$\cong_{\text{SB}} \sim \text{Hom}_{\text{cpt}}$ .

## Upper bound of $E_G^X$

Theorem (Gao-Kechris, 2003)

Let  $\cong_{\text{PM}}$  be the isometric isomorphism relation between Polish metric spaces. Then  $E_G^X \leq_B \cong_{\text{PM}}$  for every Polish  $G$ -space  $X$ .

Theorem (Melleray-Weaver, 2007)

$\cong_{\text{PM}} \sim \cong_{\text{SB}}$ .

Theorem (Zielinski, 2015)

$\cong_{\text{SB}} \sim \text{Hom}_{\text{cpt}}$ .

# Upper bound of $E_G^X$

## Theorem (Gao-Kechris, 2003)


Let  $\cong_{\text{PM}}$  be the isometric isomorphism relation between Polish metric spaces. Then  $E_G^X \leq_B \cong_{\text{PM}}$  for every Polish  $G$ -space  $X$ .

## Theorem (Melleray-Weaver, 2007)

$\cong_{\text{PM}} \sim \cong_{\text{SB}}$ .

## Theorem (Zielinski, 2015)

$\cong_{\text{SB}} \sim \text{Hom}_{\text{cpt}}$ .





# Outline

- 1 Equivalence relations and invariants
- 2 Dichotomy theorems
- 3 Orbit equivalence relations
- 4  $\Sigma_1^1$  equivalence relations

# $\Sigma_1^1$ equivalence relations

Theorem (Kechris-Louveau, 1997)

$E_1 \not\leq_B E_G^X$  for any Polish  $G$ -space  $X$ .

## Upper bound of $\Sigma_1^1$ equivalence relations

- 1  $\cong_{\text{SB}}^{\text{L}}$ : separable Banach spaces, linear isomorphism;
- 2  $\text{Hom}_{\text{SB}}^{\text{Lip}}$ : separable Banach spaces, Lipschitz isomorphism;
- 3  $\text{Hom}_{\text{PM}}^{\text{U}}$ : Polish metric spaces, uniform homeomorphism ;
- 4  $\cong_{\text{AG}}$ : abelian Polish groups, topological isomorphism.

Theorem (Ferenczi-Louveau-Rosendal, 2009)

For any  $\Sigma_1^1$  equivalence relation  $E$ , we have

$$E \leq_B \cong_{\text{SB}}^{\text{L}} \sim_B \text{Hom}_{\text{SB}}^{\text{Lip}} \sim_B \text{Hom}_{\text{PM}}^{\text{U}} \sim_B \cong_{\text{AG}} .$$

## Upper bound of $\Sigma_1^1$ equivalence relations

- 1  $\cong_{\text{SB}}^{\text{L}}$ : separable Banach spaces, linear isomorphism;
- 2  $\text{Hom}_{\text{SB}}^{\text{Lip}}$ : separable Banach spaces, Lipschitz isomorphism;
- 3  $\text{Hom}_{\text{PM}}^{\text{U}}$ : Polish metric spaces, uniform homeomorphism ;
- 4  $\cong_{\text{AG}}$ : abelian Polish groups, topological isomorphism.

Theorem (Ferenczi-Louveau-Rosendal, 2009)

For any  $\Sigma_1^1$  equivalence relation  $E$ , we have

$$E \leq_B \cong_{\text{SB}}^{\text{L}} \sim_B \text{Hom}_{\text{SB}}^{\text{Lip}} \sim_B \text{Hom}_{\text{PM}}^{\text{U}} \sim_B \cong_{\text{AG}} .$$

## Upper bound of $\Sigma_1^1$ equivalence relations

- 1  $\cong_{\text{SB}}^{\text{L}}$ : separable Banach spaces, linear isomorphism;
- 2  $\text{Hom}_{\text{SB}}^{\text{Lip}}$ : separable Banach spaces, Lipschitz isomorphism;
- 3  $\text{Hom}_{\text{PM}}^{\text{U}}$ : Polish metric spaces, uniform homeomorphism ;
- 4  $\cong_{\text{AG}}$ : abelian Polish groups, topological isomorphism.

Theorem (Ferenczi-Louveau-Rosendal, 2009)

For any  $\Sigma_1^1$  equivalence relation  $E$ , we have

$$E \leq_B \cong_{\text{SB}}^{\text{L}} \sim_B \text{Hom}_{\text{SB}}^{\text{Lip}} \sim_B \text{Hom}_{\text{PM}}^{\text{U}} \sim_B \cong_{\text{AG}} .$$

## Upper bound of $\Sigma_1^1$ equivalence relations

- 1  $\cong_{\text{SB}}^{\text{L}}$ : separable Banach spaces, linear isomorphism;
- 2  $\text{Hom}_{\text{SB}}^{\text{Lip}}$ : separable Banach spaces, Lipschitz isomorphism;
- 3  $\text{Hom}_{\text{PM}}^{\text{U}}$ : Polish metric spaces, uniform homeomorphism ;
- 4  $\cong_{\text{AG}}$ : abelian Polish groups, topological isomorphism.

Theorem (Ferenczi-Louveau-Rosendal, 2009)

For any  $\Sigma_1^1$  equivalence relation  $E$ , we have

$$E \leq_B \cong_{\text{SB}}^{\text{L}} \sim_B \text{Hom}_{\text{SB}}^{\text{Lip}} \sim_B \text{Hom}_{\text{PM}}^{\text{U}} \sim_B \cong_{\text{AG}} .$$


## Upper bound of $\Sigma_1^1$ equivalence relations

- 1  $\cong_{\text{SB}}^{\text{L}}$ : separable Banach spaces, linear isomorphism;
- 2  $\text{Hom}_{\text{SB}}^{\text{Lip}}$ : separable Banach spaces, Lipschitz isomorphism;
- 3  $\text{Hom}_{\text{PM}}^{\text{U}}$ : Polish metric spaces, uniform homeomorphism ;
- 4  $\cong_{\text{AG}}$ : abelian Polish groups, topological isomorphism.

Theorem (Ferenczi-Louveau-Rosendal, 2009)

For any  $\Sigma_1^1$  equivalence relation  $E$ , we have

$$E \leq_B \cong_{\text{SB}}^{\text{L}} \sim_B \text{Hom}_{\text{SB}}^{\text{Lip}} \sim_B \text{Hom}_{\text{PM}}^{\text{U}} \sim_B \cong_{\text{AG}} .$$





The end

Thank you!