

Alternative set-theoretic foundations of mathematics?

Yang, Ruizhi

joint work with Joel D. Hamkins

2018 Chinese Mathematical Logic Conference

May 11 - May 13, 2018

Qiannan Normal University for Nationalities

Alternative set-theoretic foundations of mathematics?

Yang, Ruizhi

joint work with Joel D. Hamkins

2018 Chinese Mathematical Logic Conference

May 11 - May 13, 2018

Qiannan Normal University for Nationalities

Definition

Let \mathfrak{A} and \mathfrak{B} be structures (not necessarily for the same first-order language). \mathfrak{A} is a **reduct** of \mathfrak{B} if its domain, relations and functions are definable in \mathfrak{B} .

Clearly, the theory of a reduct is interpretable in the theory of the original structure.

Set theory is notorious for its changeability by forcing. It is reasonable to trade-off between expressive power and solidity (effective completeness) by looking at some reducts of a set theory model.

Example (restrict the universe)

- hereditarily finite sets (HF)
- hereditarily countable sets (HC) or $L(\mathbb{R})$
- Ultimate-L

An alternative idea to reduce the expressive power is to restrict the relation “ \in ” but leave the universe as it is.

Question (Hamkins and Kikuchi 2016)

Can **set-theoretic mereology** serves as foundation of mathematics?

Set-theoretic mereology

In contrast to set theory based on the membership relation \in , **set-theoretic mereology** is the theory of the parthood relation \subset between sets.

Question

Is parthood relation **\in -complete** in the sense that \in is definable in its reduct (V, \subset) ?

Set-theoretic mereology

Fact

(V, \subset) is not \in -complete

Proof sketch: Let $\theta : V \rightarrow V$ be a nontrivial permutation.

Define

$$\tau(a) = \{\theta(b) \mid b \in a\}$$

Then $\tau : V \rightarrow V$ is a non-trivial automorphism on (V, \subset)

Set-theoretic mereology

Theorem (Hamkins and Kikuchi 2016)

Set-theoretic mereology, namely the theory of (V, \subset) is precisely the theory of an atomic unbounded relatively complemented distributive lattice. This finitely axiomatizable theory is complete and decidable.

Gödel's Thesis

A complete decidable theory can not serve as a foundation of mathematics.

Expanding the subset relation

Observation (Hamkins and Kikuchi 2016)

$(V, \subset, \{x\})$ is \in -complete

$$a \in b \Leftrightarrow \{a\} \subset b$$

Expanding the subset relation

Definition

- $a \subset^* b$ iff $a - b$ is finite
- $|a| = \infty$ iff there is a surjection from a proper subset of a to a itself

Observation

The theory of (V, \subset, \subset^*) is mutually interpretable with the theory of $(V, \subset, |\cdot| = \infty)$. The latter has a complete decidable theory.

Expanding the subset relation

We add the the unary union \bigcup or the unary intersection \bigcap instead.

Observation

Both (V, \subset, \bigcup) and (V, \subset, \bigcap) are \in -complete

$$\begin{aligned}y = \{x\} &\Leftrightarrow \bigcup y = x \wedge |y| = 1 \\ &\Leftrightarrow \bigcap y = x \wedge |y| = 1\end{aligned}$$

Expanding the subset relation

Observation

(V, \cap, \cap) , (V, \cup, \cup) , and (V, \cap, \cup) are all \in -complete

$$x \subset y \Leftrightarrow \exists z (\cap z = x \wedge \cup z = y)$$

Expanding the subset relation

Observation

(V, \cap, \cap) , (V, \cup, \cup) , and (V, \cap, \cup) are all \in -complete

$$x \subset y \Leftrightarrow \exists z (\cap z = x \wedge \cup z = y)$$

Question

What about the **unary intersection** \cap or the **unary union** \cup on themselves?

The reduct (V, \cap) is just a proper-class-branching “tree” of height \aleph_0 if (V, \in) is well-founded

Theorem

There is a complete axiomatization of the theory of (V, \cup) .

The unary union structure

There are exactly two **covers** of \emptyset :

$$\bigcup \emptyset = \emptyset$$

$$\bigcup \{\emptyset\} = \emptyset$$

Note that \emptyset is the only finite set x such that $\bigcup x = x$.

The unary union structure

There are exactly two covers of $\{\emptyset\}$:

$$\bigcup\{\{\emptyset\}\} = \{\emptyset\}$$

$$\bigcup\{\emptyset, \{\emptyset\}\} = \{\emptyset\}$$

Note that it is the case for any singleton $\{a\}$.

The unary union structure

There are exactly two covers of $\{a\}$:

$$\cup\{\{a\}\} = \{a\}$$

$$\cup\{\emptyset, \{a\}\} = \{a\}$$

Note that it is the case for any singleton $\{a\}$.

The unary union structure

How many covers are there for a set of size **two** $\{a, b\}$?

$\{\{a, b\}\}; \{\emptyset, \{a, b\}\}; \{\{a\}, \{a, b\}\}; \{\{b\}, \{a, b\}\}; \{\{a\}, \{b\}\};$

$\{\emptyset, \{a\}, \{a, b\}\}; \{\emptyset, \{b\}, \{a, b\}\}, \{\emptyset, \{a\}, \{b\}\}; \{\{a\}, \{b\}, \{a, b\}\};$

$\{\emptyset, \{a\}, \{b\}, \{a, b\}\}.$

The unary union structure

How many covers are there for a set of size **two** $\{a, b\}$?

$\{\{a, b\}\}; \{\emptyset, \{a, b\}\}; \{\{a\}, \{a, b\}\}; \{\{b\}, \{a, b\}\}; \{\{a\}, \{b\}\};$

$\{\emptyset, \{a\}, \{a, b\}\}; \{\emptyset, \{b\}, \{a, b\}\}, \{\emptyset, \{a\}, \{b\}\}; \{\{a\}, \{b\}, \{a, b\}\};$

$\{\emptyset, \{a\}, \{b\}, \{a, b\}\}.$

The unary union structure

How many covers are there for a set of size **three** $\{a, b, c\}$?

There are **218** covers of a set containing three elements!

- The number of covers of **4** is **64594**.
- The number of covers of **5** is **4294642034**.

.....

The unary union structure

How many covers are there for a set of size **three** $\{a, b, c\}$?

There are **218** covers of a set containing three elements!

- The number of covers of **4** is **64594**.
- The number of covers of **5** is **4294642034**.

.....

The unary union structure

How many covers are there for a set of size **three** $\{a, b, c\}$?

There are **218** covers of a set containing three elements!

- The number of covers of **4** is **64594**.
- The number of covers of **5** is **4294642034**.

.....

The unary union structure

Fact

There is a recursive function C , given a finite set

$A = \{a_1, \dots, a_n\}$, there are exactly $C(n)$ many covers of A .

$$C(n) = 2^{2^n} - \left(\sum_{i=0}^{n-1} \binom{n}{i} \cdot C(i) \right).$$

$$C(n) = \sum_{i=0}^n (-1)^{n-i} \cdot \binom{n}{i} \cdot 2^{2^i}.$$

The unary union structure

Fact

There is a recursive binary function c such that, given a finite set A of size n , there are $c(n, m)$ many covers of A of size m .

$$c(n, m) = \binom{2^n}{m} - \left(\sum_{i=0}^{n-1} \binom{n}{i} \cdot c(i, m) \right).$$

$$c(n, m) = \sum_{i=0}^n (-1)^{n-i} \cdot \binom{n}{i} \cdot \binom{2^i}{m}.$$

The unary union structure

The infinite case

Let κ be infinite, $\lambda \geq 2$. There are $C(\kappa) = 2^{2^\kappa}$ many covers of κ , among them, there are $c(\kappa, \lambda) = [2^\kappa]^\lambda$ many covers of cardinality λ .

The unary union structure

Some definable sets in (V, \cup) :

- $x = \emptyset \Leftrightarrow \bigcup x = x \wedge \exists^{=2} y \bigcup y = x$
- $x = \{\emptyset\} \Leftrightarrow \bigcup x = \emptyset \wedge x \neq \emptyset$
- $x = \{\{\emptyset\}\} \Leftrightarrow \bigcup x = \{\emptyset\} \wedge \exists^{=2} y \bigcup y = x$
- $x = \{\emptyset, \{\emptyset\}\} \Leftrightarrow \bigcup x = \{\emptyset\} \wedge \exists^{=10} y \bigcup y = x$

The unary union structure

Observation

For each $1 \leq k < \omega$ there are infinitely many sets $x \in V_{\omega+1}$ such that x have exactly k many \cup -successors in $V_{\omega+1}$ (it follows $\bigcup^k x = x$) constituting a k -loop.

Example

- $\{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \dots\}$
- $\{\emptyset, \{\emptyset\}, \{\{\emptyset\}, \emptyset\}, \{\{\{\emptyset\}, \emptyset\}\}, \dots\}$
- $\{\emptyset, \{\{\emptyset\}\}, \{\{\{\{\emptyset\}\}, \emptyset\}\}, \dots\}$

Axioms of union

Definition

$$D0 \quad |x| = 0 \leftrightarrow \bigcup x = x \wedge \exists^{=2} y \bigcup y = x$$

$$Dn \quad |x| = n \leftrightarrow \bigcup x \neq x \wedge \exists^{=C(n)} y \bigcup y = x$$

$$E1 \quad |x| \geq 0 \leftrightarrow x = x$$

$$En \quad |x| \geq n \leftrightarrow \exists^{\geq C(n)} y \bigcup y = x \wedge y \neq x$$

Note: $|x| = 0 \vee \dots \vee |x| = n - 1 \vee |x| \geq n$ is not valid

Axioms of union

U1

$$\exists^=1 x |x| = 0$$

U2 For $n, k \in \omega$,

$$\exists^{\geq n} x (x = \bigcup^k x \wedge \bigwedge_{1 \leq k < k} x \neq \bigcup^l x)$$

U3 For $n \geq 1$,

$$\forall x \left(\bigvee_{i=0}^{n-1} |x| = i \vee |x| \geq n \right)$$

Axioms of union

U4 For $n, k \geq 1$,

$$\forall x (|x| = n \rightarrow x \neq \bigcup^k x)$$

U5 For $n, m \in \omega$,

$$\forall x (|x| \geq n \rightarrow \exists^{\geq c(n,m)} y (\bigcup y = x \wedge |y| = m))$$

U6

$$\forall x \exists^=1 y (\bigcup y = x \wedge |y| = 1)$$

Quantifier elimination

Let $\mathcal{L}_U^* = \{\cup, \emptyset, |x| = n, |x| \geq n\}$. Let Λ_U^* be constituted by **U1** - **U6**, **Dn** s, **En** s, and $|\emptyset| = 0$

Lemma

For every \mathcal{L}_U^* -formula θ , there is a quantifier-free \mathcal{L}_U^* -formula ψ such that

$$\Lambda_U^* \vdash \theta \leftrightarrow \psi$$

Corollary

$\Lambda_U = \mathbf{U1} - \mathbf{U6}$ is complete

More reducts

Theorem

(V, \cap) has a decidable theory:

I1 $\exists^{=1} x \cap x = x$

I2 $\forall x (\cap^k x = x \rightarrow \cap x = x)$ (for $k > 1$)

I3 $\forall x \exists^{\geq n} y \cap y = x$ (for $n < \omega$)

More reducts

Theorem

(V, \cup, P) has a decidable theory: U1 - U6 together with

$$P1 \quad \forall x (|Px| \neq n \rightarrow \bigvee_{m \leq \lfloor \log n \rfloor} |Px| = 2^m) \text{ (for } n < \omega)$$

$$P2 \quad \forall x (|x| = n \leftrightarrow |Px| = 2^n) \text{ (for } n < \omega)$$

$$P3 \quad \forall x \cup Px = x;$$

$$P4 \quad \forall x (P^l x \neq \bigcup^k x) \text{ (for } l > 1 \text{ and } k < \omega)$$

Summery

ϵ -complete:

$(V, \subset, \{x\})$, (V, \subset, \cup) , (V, \subset, \cap) , (V, \cup, \cap) , etc.

completely axiomatizable:

$(V, \subset, |\mathcal{X}| = \infty)$, (V, \cup) , (V, \cap) , (V, \cup, P) , etc.

Theorem

The reduct (V, \subset, P) is rigid, namely every nontrivial automorphism on (V, \subset) does not preserve P

Question

Is (V, \subset, P) ϵ -complete?

Thank you!